FOR USE IN ANIMALS ONLY

INDICATIONS: For use as a supplemental nutritive source of vitamins A and D in cattle. **CAUTION:** Do not inject into meat animals within 60 days of marketing. Administration of this product to well nourished animals may cause hypervitaminosis D which may result in hypercalcemia. This condition may be harmful to the animal.

DOSAGE AND ADMINISTRATION:

Inject intramuscularly or subcutaneously preferably in the neck area using aseptic technique.

Calves - 1/2 to 1 mL Yearling Cattle - 1 to 2 mL Adult Cattle - 2 to 4 mL These suggested dosages may be repeated after 60 days, if necessary.

Manufactured for: **DURVET, INC.** Blue Springs, Missouri 64014 www.durvet.com





INJECTABLE STERILE SOLUTION

For use as supplemental nutritive source of vitamins A and D in cattle.

NET CONTENTS:

500 mL

(16.9 FL. OZ.)

KEEP OUT OF REACH OF CHILDREN

Each mL of sterile solution contains:

OTTL	INGILEDIEN IO.	
Vitamin	Α	500,000 IU
Vitamin	D ₃	75,000 IU

INACTIVE INGREDIENTS: Emulsifiable base with vitamin E (antioxidant), N-methylpyrrolidone, polyoxyethylated (30) castor oil, propylene glycol dicaprylate, polyoxyethylene (20) sorbitan monooleate and benzyl alcohol 2%.

Store at controlled room temperature between 15° and 30°C (59°-86°F).

Protect from light and excessive

heat.

V-0332-06 REV 02-24

			_
			_
. =	_	_	_
-			M
-			
			· +
-			~
			_
			· ·
			· ^ I
			<u></u>
			_
- 2			_
- 22			· ·
-			
			∞
			ഹ
			_
			~