

FOR USE IN ANIMALS ONLY

INDICATIONS:

For use as an aid in the treatment of acetonemia (Ketosis) in cattle.

CAUTION: Intravenous administration must be done slowly and made under strict asepsis. Solution should be warmed to body temperature prior to administration.

This is a single dose container. This product contains no preservative. After a quantity has been withdrawn for injection, the remainder should be discarded. Do not administer intraperitoneally.



ISS19XB04

Manufactured for:
DURVET, INC.

Blue Springs, Missouri 64014
www.durvet.com

NDC 30798-127-17



Dextrose 50% SOLUTION

Dura-Ster TS
TERMINALLY STERILIZED



For use as an aid in the treatment of acetonemia (Ketosis) in cattle.

NET CONTENTS:
500 mL
(16.9 fl. oz.)

KEEP OUT OF REACH OF CHILDREN

EACH 100 mL CONTAINS:

Dextrose •H₂O..... 50 gms

DOSAGE AND ADMINISTRATION:

For intravenous administration only.

Cattle: 100 to 500 mL depending on size and condition. Treatment may be repeated in several hours or on successive days as needed.

Store between 15°C and 30°C (59°F-86°F)

TAKE TIME  OBSERVE LABEL DIRECTIONS



Terminally Sterilized to provide a greater degree of sterilization.