

**FOR USE IN ANIMALS ONLY**  
**INDICATIONS:**

Dextrose 50% Solution is indicated for use as an aid in the treatment of uncomplicated primary ketosis in cattle.

**DOSAGE & ADMINISTRATION:**

The usual dose is 50 mL per 100 lbs of body weight. It should be administered intravenously only. Dosage may be repeated in 8 to 10 hours or on successive days if necessary.

**CAUTION:**

It should be warmed to body temperature and administered slowly. This product contains no preservatives. Do not use if solution is not clear. Entire contents should be used upon opening. Discard any unused portion.



NDC 30798-856-17



**Dextrose 50%**  
**SOLUTION**

**Sterile Solution**



For use as aid in the treatment of uncomplicated primary ketosis in cattle.

**NET CONTENTS:**  
**500 mL**  
**(16.91 FL. OZ.)**

**KEEP OUT OF REACH OF CHILDREN**

**ACTIVE INGREDIENTS:**

Dextrose • H<sub>2</sub>O ..... 50% w/v  
Water for Injection ..... q.s.

**Store between 15°C - 30°C (59°F - 86°F).**

TAKE TIME



OBSERVE LABEL DIRECTIONS

ISS24XB11 RMS 92-324

Manufactured for:  
**DURVET, INC.**  
Blue Springs, Missouri 64014  
[www.durvet.com](http://www.durvet.com)



Lot No.

Exp. Date