



# Chick Starter

## TYPE C MEDICATED FEED

### for replacement chickens

with essential oils



For development of active immunity

## CRUMBLES

For development of active immunity to coccidiosis under conditions of slight to severe exposure to coccidiosis in replacement chickens up to 8 weeks of age.

### ACTIVE DRUG INGREDIENT

Amprolium.....0.0125% (113.5 g/ton)

See back panel for instructions for use and warning statements.



NET WT:  
10 LB (4.54 KG)



# Chick Starter

## TYPE C MEDICATED FEED

replacement chickens

**For development of active immunity to coccidiosis under conditions of slight to severe exposure to coccidiosis in replacement chickens up to 8 weeks of age.**

### ACTIVE DRUG INGREDIENT

Amprolium.....0.0125% (113.5 g/ton)

### GUARANTEED ANALYSIS

Crude Protein, min.....19.5%  
Lysine, min ..... 1.00%  
Methionine, min ..... 0.40%  
Crude Fat, min ..... 2.75%  
Crude Fiber, max ..... 3.50%  
Calcium, min ..... 0.75%  
Calcium, max ..... 1.25%  
Phosphorus, min ..... 0.65%  
Salt, min ..... 0.25%  
Salt, max .....0.45%

### INGREDIENTS

Grain products, plant protein products, monocalcium phosphate, calcium carbonate, sodium chloride, choline chloride, dl-methionine, Lysine, Vitamin A supplement, Vitamin D3 supplement, Vitamin E supplement, Niacin supplement, Calcium Pantothenate, Menadione dimethylpyrimidinol bisulfite, Thiamine Mononitrate, Riboflavin, Pyridoxine Hydrochloride, Vitamin B12 supplement, Biotin, Folic Acid, Iron Sulfate, Copper Sulfate, Basic Copper Chloride, Zinc Hydroxychloride, Zinc Sulfate, Manganous Oxide, Manganese Hydroxychloride, Sodium Selenite, Ethylenediamine dihydriodide, Essential Oils of Oregano (Carvacrol), Red Pepper (Capsicum), Cinnamon (Cinnamaldehyde), Thyme, and Rosemary Extract and Vegetable Oil.

### FEEDING DIRECTIONS

Feed continuously as the sole ration to replacement chickens up to 8 weeks of age. At 6 weeks of age, one could begin feeding a grower ration containing amprolium up to 8 weeks of age.

### CAUTION

Use as the sole source of amprolium. Do not change the litter while giving this feed unless absolutely necessary. If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis, and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of flocks and prompt treatment will be required. In replacement flocks grower must expect that excessive exposure to one or more species may overwhelm the drug in some flocks and prompt treatment will be required. Do not use in feeds containing bentonite.

### WARNING

**Not for human use. Keep this and all drugs out of reach of children.**

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report adverse effects, obtain an SDS or for assistance, contact Durvet at 1-800-821-5570.



Manufactured for Durvet, Inc., Blue Springs, MO 64014  
www.durvet.com | 1-800-821-5570

**NET WT: 10 LB (4.54 KG)**