

## Feline Leukemia-**Rhinotracheitis-**Calici-Panleukopenia Vaccine

Modified Live & Killed Virus

## Focus Cat Vax 3

Package contains a one dose vial of vaccine. a one mL vial of diluent, and one disposable syringe

PERISHABLE PRODUCT • KEEP REFRIGERATED

Indications: This product has been shown to be effective for the vaccination of healthy cats 8 weeks of age or older against feline leukemia, rhinotracheitis, calici and panleukopenia viruses. The duration of immunity for this product has not been established. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.

Directions and dosage: Open syringe by twisting or tapping the cap against a hard surface to break the heat weld. Aseptically rehydrate the vaccine cake with 1mL of FeLV sterile diluent supplied. SHAKE WELL, Withdraw entire contents into the syringe. Administer one 1 mL dose subcutaneously using aseptic technique: lift the loose skin behind the neck or behind the right hind leg (see illustration) and insert needle. Inject entire contents of syringe. Do not inject directly into blood vessel. IMPORTANT NOTE: Before injecting vaccine, pull back slightly on syringe plunger. If blood enters the syringe freely, choose another injection site. Healthy cats should receive

two doses, 3 to 4 weeks apart, except that if the animal is less than 16 weeks of age, the final dose should be given no earlier than 16 weeks of age. The presence of maternal antibody is known to interfere with the development of active immunity in cats and additional boosters will be required in most voung animals. Historically, annual revaccination with a single dose has been recommended for this product. The need for this booster has not been established. For advice on revaccination frequency and annual booster vaccinations. consultation with a veterinarian is recommended.

Precautions: DO NOT VACCINATE PREGNANT QUEENS. Store in the dark at 35-46°F (2-8°C), DO NOT FREEZE, Do not mix with other products, except as specified on this label. In case of anaphylactoid reaction, administer epinephrine. In case of human exposure, contact a physician. Inactivate unused contents before disposal. Thimerosal, neomycin, polymyxin B, and amphotericin B added as preservatives.





VaccinateYourPets.com

Produced by:

Elanco US Inc. Fort Dodge, IA 50501 USA VLN/PCN 196/15B9.21

Distributed by: Peak Marketing Blue Springs MO 64014 USA Phone: 1-800-821-5570

Feline Focus® & Cat Vax 3® are registered trademarks of Durvet. Inc. ISS20XB08

3G102784X

Step 1 Transfer liquid contents of diluent into vial of dried vaccine Step 2

Shake well

Step 3 Withdraw mixture into syringe

Step 4 To administer subcutaneously, gently grasp and elevate a fold of loose skin

Insert the needle into the loose tissue under the skin



žer.: