

Lamb & Kid Colostrum Replacer

RECLOSABLE BAG

MEASURING SCOOP INCLUDED
(0.5 oz (0.014 KG))



Dura-Start

LAMB & KID

Colostrum Replacer

8 grams of Globulin Protein per 2 oz. of powder



A **D₃** Formulated with essential vitamins and minerals

Convenient, easy mixing for reliable nutritional consistency

NET WT. 10 oz. (283.5 g)



01-2213-0010 2896545
ISS24XB02
www.durvet.com
Blue Springs, MO 64014
100 S.E. Magellan Drive
Durvet, Inc.

Manufactured for

STORAGE

Store dry powder in a tightly closed container in a cool, dry place.
Pyridoxine Hydrochloride, Ethylenediamine Dihydrochloride, Riboflavin Supplement, Cobalt Sulfate, Thiamine Mononitrate, Biotin, Calcium Lactate, Vitamin B12 Supplement, Menadione Sodium Bisulfite Complex (Vitamin K), Folic Acid, Vitamin A Supplement, Vitamin E Supplement, Sodium Sulfate, Zinc Sulfate, Manganese Sulfate, Selenium, Ferrous Sulfate, Vitamin D3 Supplement, Calcium Acetate, Choline Chloride, Vitamin E Supplement, Sodium L-Lysine, Natural and Artificial Flavor, DL-Methionine, Ascorbic Acid, Calcium Phosphate, Calcium Carbonate, Calcium Silicate, Dextrose, Coconut Oil (preserved with citric acid), Lecithin, Animal Fat (preserved with BHA and BHT), Dried Whey, Dried Whey Protein Concentrate, Dried Bovine Colostrum, INGREDIENTS

Guaranteed Analysis
Crude Protein, Min. 35.00%
Crude Fat, Min. 25.00%
Crude Fiber, Max. 0.10%
Calcium, Min. 0.75%
Calcium, Max. 1.25%
Phosphorus, Min. 0.70%
Sodium, Max. 1.00%
Selenium, Min. 0.30 ppm
Vitamin A, Min. 20,000 IU/lb
Vitamin D3, Min. 5,000 IU/lb
Vitamin E, Min. 150 IU/lb
Globulin Protein, Min. 14.30%

INGREDIENTS

CAUTION
Follow label directions. The addition to feed of higher levels of this premix containing selenium is not permitted.
Not for human consumption. Wash hands after feeding/handling animals. Wash feed/water containers after use.
MIXING DIRECTIONS: Mix 2 ounces (~4 scoops) of Dura-Start Lamb & Kid into 7 ounces of warm water (110-120°F). Feed solution at body temperature (~104°F). See the feeding chart below for mixing directions.
Total # of Scoops to Lamb or Kid Birthweight
Total # of Scoops to Replace Maternal Colostrum
Total # of Scoops to Supplement Maternal Colostrum
of Feedings*

4 lb	4	1	2
8 lb	8	2	2-3
12 lb	12	3	2-3
14 lb	16	4	2-3

* Plan 2 to 3 feedings of mixed solution within the first 8 hours.
** Enclosed scoop holds approximately 0.5 ounces of powder.
Weigh powder for best accuracy.
Refrigerate unused colostrum supplement solution promptly after mixing. Reheat gently over warm water and mix thoroughly before feeding. Do not microwave or freeze the solution. Discard any unused refrigerated colostrum supplement solution after 8 hours.
DIRECTIONS FOR USE: (See chart above for Mixing Instructions)
When Fed as a Replacer: Feed 7 fluid ounces of mixed solution per 4 lb of bodyweight. Split this mixed solution into 2-3 feedings over the first 8 hours of life.
When Fed as a Supplement: Feed 3.5 fluid ounces of mixed solution per 4 lb of bodyweight. The mixed solution can be added to maternal colostrum if there is not enough quantity or if quality of the maternal colostrum is unknown as may be the case in multiple births. Lambs and kids also benefit from feeding supplemental colostrum past day one.
Offer fresh water at all times.